



# hDGAT2

利用同源重组，将小鼠Dgat2基因进行人源化修饰。

<b>Nomenclature</b>	C57BL/6JSm0-Dgat2 <sup>tm1(hDGAT2)Smoc</sup>
<b>Cat. NO.</b>	NM-HU-233515
<b>Strain State</b>	Repository Live

## Gene Summary


<b>Gene Symbol</b> Dgat2	<b>Synonyms</b>	ARAT; DGAT-2; 0610010B06Rik
	<b>NCBI ID</b>	<a href="#">67800</a>
	<b>MGI ID</b>	<a href="#">1915050</a>
	<b>Ensembl ID</b>	<a href="#">ENSMUSG00000030747</a>
	<b>Human Ortholog</b>	DGAT2

## Model Description

利用同源重组，将小鼠Dgat2基因进行人源化修饰。

\*Literature published using this strain should indicate: hDGAT2 mice (Cat. NO. NM-HU-233515) were purchased from Shanghai Model Organisms Center, Inc..

## Validation Data




### Fig.1 Detection of DGAT2 expression in the liver and subcutaneous fat pad by RT-PCR.

Wild type: only one band at 481 bp with primers F1/R1(*mDgat2*);

Heterozygous: one band at 481 bp with primers F1/R1(*mDgat2*) and one band at 388 bp with primers F2/R2(*hDGAT2*).

Abbr. M, DNA marker; HE, heterozygous; WT, wild type.




### Fig.2 Detection of human DGAT2(A) and mouse Dgat2(B) mRNA expression in liver and WAT by qPCR.

A: Relative expression was reported as the percentage to human DGAT2 mRNA levels in the liver of hDGAT2 knockin mice (n=2, male, 9-week-old).

B: Relative expression was reported as the percentage to mouse Dgat2 mRNA levels in the liver of WT C57BL/6 mice (n=2, male, 9-week-old).

Abbr. M, DNA marker; HO, homozygous; WT, wild type.



**Fig.3 Detection of human DGAT2 mRNA knockdown in liver and gWAT by qPCR.** Female mice (8-9 weeks old) were assigned to three groups (n=2 per group). Homozygous hDGAT2 (HO) mice received a single subcutaneous injection of PBS (Groups 1), while HO mice in Group 2 were administered a single subcutaneous dose of ION224, a GalNAC-conjugated ASO drug (10 mg/kg). At 15 days post-dosing, liver and gonadal white adipose tissue (gWAT) were harvested to quantify human *DGAT2* mRNA levels. Data were normalized to mouse *GAPDH* mRNA levels. Results demonstrated that ION224 significantly reduced human *DGAT2* mRNA expression by 80% in the liver and approximately 50% in the gWAT compared to the PBS-treated HO group.

Abbr. HO, homozygous.